



## ORIGINAL ARTICLE



WILEY

# Effects of ocean climate on the length and condition of forage fish in the Gulf of Alaska

Sarah Ann Thompson<sup>1</sup> | Marisol García-Reyes<sup>1</sup> | William J. Sydeman<sup>1</sup> |  
Mayumi L. Arimitsu<sup>2</sup> | Scott A. Hatch<sup>3</sup> | John F. Piatt<sup>4</sup>

<sup>1</sup>Farallon Institute, Petaluma, California

<sup>2</sup>U.S. Geological Survey, Alaska Science Center, Juneau, Alaska

<sup>3</sup>Institute for Seabird Research and Conservation, Anchorage, Alaska

<sup>4</sup>U.S. Geological Survey, Alaska Science Center, Anchorage, Alaska

**Correspondence**

Sarah Ann Thompson, Farallon Institute, Petaluma, CA.

Email: sathompson@faralloninstitute.org

**Funding information**

North Pacific Research Board, Grant/Award Number: Project 1706; National Fish and Wildlife Foundation; U.S. Geological Survey

**Abstract**

Climatic drivers of the size and body condition of forage fish in the North Pacific are poorly known. We hypothesized that length and condition of forage fish in the Gulf of Alaska (GoA) vary in relation to ocean temperature on multiple scales. To test this hypothesis, we analyzed morphometric data for capelin (*Mallotus catervarius*) and Pacific sand lance (PSL; *Ammodytes personatus*) sampled by a seabird (*Cerorhinca monocerata*) in two regions of the GoA, 1993–2016. Based on previous studies, we predicted that capelin length and body condition (Fulton's *K*) would be negatively related to the Pacific Decadal Oscillation (PDO) and sea surface temperature (SST), whereas PSL length and condition would be positively related. Interannual variation in length and body condition was evaluated relative to seasonal values of ocean climate using regression. Forage fish length and condition varied interannually, between sampling regions, and were dependent on the size/age class of the fish sampled. As predicted, length and body condition of capelin (mostly age 1+) were negatively related to the PDO and SST. Relationships with ocean climate for PSL varied by size/age class: positive for putative age-0 fish and negative for putative age-1+ fish. We conclude that our hypothesis was supported for capelin and partially supported for PSL. This study demonstrates that ocean climate determines key morphometric characteristics of forage fish that may relate to interannual variation in the energetic value of prey, and provides an example of how seabirds can be used to obtain specimens for evaluations of potential prey quality.

**KEYWORDS**

condition, forage fish, interannual variation, ocean climate, seabird

## 1 | INTRODUCTION

Forage fish quantity and quality have important effects on the ecology and vital rates of marine top predators, including seabirds (Suryan et al., 2002), marine mammals (Trites & Donnelly, 2003), and large predatory fish (Rose & O'Driscoll, 2002). In particular, the “junk food” hypothesis describes ways in which lower food quality may negatively affect marine predator population parameters (Alverson, 1992; Rosen & Trites, 2000; reviewed by Osterblom,

Olsson, Blenckner, & Furness, 2008). While it has been well established that prey quality is important to aspects of predator success, drivers of variation in quality remain poorly understood. Only a few studies have been conducted over a long enough period of time to examine interannual or lower-frequency variation in prey quality (Carscadden, Montevecchi, Davoren, & Nakashima, 2002; Wanless, Harris, Newell, Speakman, & Daunt, 2018; Wanless, Harris, Redman, & Speakman, 2005). Developing a robust understanding of variation in prey size and energetic value is becoming more important with

increasing climate variability and unidirectional change (IPCC, 2013), which are likely to affect prey quality through related factors such as prey size and mass (Sheridan & Bickford, 2011).

Prey quality can be assessed by measurements of whole-body lipid and/or protein content, which directly reflect the energetic and nutritional value of prey (Anthony, Roby, & Turco, 2000; Payne, Johnson, & Otto, 1999; Van Pelt, Piatt, Lance, & Roby, 1997; Vollenweider, Heintz, Schaufler, & Bradshaw, 2011; von Biela et al., 2019). Spatial (Abookire & Piatt, 2005) and temporal variation (Litz et al., 2010; Montevecchi & Piatt, 1984; Robards, Anthony, Rose, & Piatt, 1999; Vollenweider et al., 2011) in lipid and protein content has been documented and in some cases has been related to the ocean conditions that alter the planktonic prey base that forage fish consume (Litz et al., 2010). Variation in energy content is primarily related to variation in lipid content and length (von Biela et al., 2019). Lipid content can also be inferred by measurements of body condition that index mass (g) scaled to size (mm; Robards, Anthony, et al., 1999; Logerwell & Schaufler, 2005). Consequently, measurements of body condition, such as the commonly used Fulton's *K* index (Ricker, 1975), can be used as a proxy for prey quality in space and time (Carscadden & Frank, 2002; Davoren & Montevecchi, 2003; Pangle & Sutton, 2005; Robards, Anthony, et al., 1999).

In this study, we derive length and a body condition index from forage fish samples obtained during studies of seabird diets in the Gulf of Alaska (GoA) taken over multiple decades (Hatch & Sanger, 1992; Piatt et al., 2018; Sydeman, Piatt, et al., 2017). This application of predator-based sampling (Mills, Laidig, Ralston, & Sydeman, 2007; Thayer et al., 2008) has been used to study basin-scale and annual-to-decadal variation in forage fish availability in this region (Piatt et al., 2018; Sydeman, Piatt, et al., 2017) and elsewhere in the world. For example, using common murre (*Uria aalge*) and Atlantic puffins (*Fratercula arctica*) as samplers in the North Sea, Wanless et al. (2005, 2018) showed that the energy content of sandeel (*Ammodytes marinus*) and sprat (*Sprattus sprattus*) prey was extraordinarily low in 2004 and that the overall quality of sandeels declined steadily over a 40-year sampling period. Similarly, in a decade-long study in the Northwest Atlantic, Davoren and Montevecchi (2003) showed that body condition of capelin sampled by murre declined through time, accompanied by a synchronous decline in the condition of murre chicks.

An important assumption in these studies is that seabirds sample prey in proportion to the abundance and/or availability of prey in the environment. Rarely are contemporaneous data collected on diet and prey availability, but in cases where this has been done, diet composition reflects prey abundance. This includes studies on the take of (a) age-0 cod (*Gadidae*) by puffins in the GoA (Hatch & Sanger, 1992), (b) age-0 rockfish (*Sebastes* spp.) by murre in the central California Current (Mills et al., 2007), (c) age-0 and age-1+ Pacific sand lance (PSL; *Ammodytes personatus*) and capelin (*Mallotus catervarius*) by puffins in the GoA and Aleutian archipelago (Piatt et al., 2018), and (d) age-0 and older anchovy (*Engraulis encrasicolus*) and sardine (*Sardinops sagax*) by Cape gannets (*Morus capensis*) in the southern Benguela ecosystem (Crawford, Sydeman, Thompson,

Sherley, & Makhado, 2019). We therefore assume that sampling in proportion to abundance also occurred in this study. Another critical assumption is that the seabirds sample prey sizes in proportion to those available in the environment.


The samples used in this study were obtained from rhinoceros auklets (*Cerorhinca monocerata*), a medium-sized (~500 g) puffin (Friesen, Baker, & Piatt, 1996) with the capacity to search and forage to 10–60 m depth (Burger, Wilson, Garnier, & Wilson, 1993; Cunningham, Elliott, Cottenie, Hatch, & Jacobs, 2018; Kuroki et al., 2003), typically 30–80 km from a breeding colony (Cunningham et al., 2018). Here, we focus on two key forage fish of the Alaskan North Pacific: capelin and Pacific sand lance. Our study region spans the eastern to central-northern GoA, an ecosystem highlighted by a rich and abundant community of marine mammals, fish, and seabirds, as well as valuable fisheries for salmon, cod, and halibut. Biological productivity and the controls of energy flow from lower to upper trophic levels in this region are related to water movement of the Alaska Gyre, winds and upwelling/downwelling, ocean temperatures, and stratification (Brodeur & Ware, 1992; Stabeno et al., 2004, 2016; Waite & Mueter, 2013).

Our focal forage fish species have different habitat preferences within the cool subarctic/boreal waters in which both species reside: PSL prefer warmer, coastal waters (Abookire & Piatt, 2005; Piatt et al., 2018; Sydeman, Piatt, et al., 2017), whereas capelin are found in deeper, colder shelf waters (Abookire & Piatt, 2005; Logerwell, Stabeno, Wilson, & Hollowed, 2007). In this study, we test the hypothesis that forage fish body condition varies in relation to large-scale to regional ocean climate parameters that regulate ecosystem productivity, primarily temperature. We predict that variability in ocean climate affects the size (proxied by length) and body condition of these species in opposing directions, with temperature relating positively to PSL size and body condition and negatively to capelin size and body condition. Past research has primarily focused on species composition and abundance in predator diet studies, but some life history events, such as breeding failures, cannot be explained by diet composition. This study is significant because understanding environmental effects on prey quality is useful for predicting impacts on predator communities and fisheries in the face of continued climate variability.

## 2 | METHODS

### 2.1 | Forage fish data

Other than for herring (*Clupea pallasii*), annual abundance estimates are not available for any forage fish species in Alaska. From the 1970s to the present, however, summer studies of puffin food habits have been conducted as part of the Alaska Maritime National Wildlife Refuge (AMNWR) seabird monitoring program as well as the sampling programs conducted by the U.S. Geological Survey Alaska Science Center and the Institute for Seabird Research and Conservation (Byrd, 2007; Hatch, 2013; Schoen et al., 2018). Field research sampling methods are detailed by Hatch and Sanger (1992) and AMNWR (2018), and the database housing the data collected



**FIGURE 1** Map showing the Gulf of Alaska with the two study sites Middleton Island (59.438°N, 146.327°W) and St. Lazaria Island (56.987°N, 135.705°W). Colors indicate average SST from April to June (°C). The boxes around each of the sites indicate the bounds for determining local SST; see Methods for details [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Species	Location	N years	N fish	Fulton's K		
				Mean	SD	CV
Capelin	Middleton	12	2,351	0.478	0.172	0.359
	St. Lazaria	21	1,688	0.507	0.090	0.178
Total			4,039			
Pacific sand lance						
Age 0 (<100 mm)	Middleton	12	2,649	0.269	0.081	0.300
	St. Lazaria	16	924	0.311	0.280	0.901
Age 1+ (≥100 mm)	Middleton	13	1,551	0.322	0.139	0.431
	St. Lazaria	18	1,315	0.329	0.056	0.171
Total			6,439			

**TABLE 1** Sample sizes and summary statistics (mean, standard deviation, and coefficient of variation; calculated based on the number of fish) of condition indices for capelin and Pacific sand lance by site, 1993–2016. See text for details on Fulton's K calculations

from these monitoring programs, the Alaska Puffin Diet Database, is described in more detail elsewhere (Piatt et al., 2018; Sydeman, Piatt, et al., 2017). In brief, researchers collected the fish that were caught as prey by adult puffins and brought to colony burrows that contain their chicks. These fish were identified and measured in the field or frozen for measurement in the laboratory; the vast majority was measured in the field. We used data from two sites in the GoA: Middleton Island (59.438°N, 146.327°W), in the northern GoA region, and St. Lazaria Island (56.987°N, 135.705°W), in the eastern GoA region (Figure 1). These sites were the most suitable for use in our study because of their locations in the GoA and because they had the most complete time series of the sites in the seabird monitoring program.

In this study, we use measurement data for capelin and PSL sampled by rhinoceros auklets between June 18 and August 26 each

year, from 1993 to 2016. These fish are among the most important forage fish in the GoA, and there were large sample sizes available for them at our study sites (Table 1). The data were filtered for records with length (mm) and mass (g) measurements on whole fish. Records with no mass or length measurement were omitted from analysis. Our database had more records that included total length than fork length. Therefore, to use all available data, we converted fork length measurements to total length with the following regressions (established from data from fish samples with both measurements): capelin total length =  $4.944 + (1.012 \times \text{fork length})$ ,  $R^2 = 0.95$ ,  $n = 1685$ ; and PSL total length =  $0.880 + (1.023 \times \text{fork length})$ ,  $R^2 = 0.99$ ,  $n = 2,298$ . For further analysis, we used the measured or predicted total length data. We assume length to be an indicator of age classes under average environmental conditions; under anomalous circumstances, some fish may be smaller (von Biela et al., 2019), and may not fall

precisely into these age classifications. Nevertheless, under this assumption, length–frequency distributions were used to identify size/age classes for each species.

We calculated Fulton's *K* index of body condition (Ricker, 1975) using the following equation:

$$K = \left( \frac{\text{mass}}{\text{length}^3} \right) \times 100$$

We chose this condition index because it is commonly used in fisheries science (Nash, Valencia, & Geffen, 2006) and was previously found to be a suitable descriptor of variation in body condition in capelin and PSL (Baker et al., 2019; Carscadden & Frank, 2002). Fulton's *K* represents variation primarily in the mass of fish, which can be variable even at a constant length (see Tables S1 and S2).

To obtain robust time series, we omitted "site-years" when <10 individual fish were measured (by species; see Table 1 for final number of years and fish). Trends in the annual *K* were described with linear and quadratic regression, weighted by sample size in order to adjust for years with few specimens. While a few statistically significant trends were found (Tables S3 and S4), these data were not detrended for analysis because most of the variables had no trends. We employed analysis of covariance (ANCOVA) to test for inter-annual variability in *K*, adjusted for sampling date. Additionally, we used an unpaired two-sided *t* test to test for differences between the mean length and condition at sites within species (Middleton and St. Lazaria for each of capelin, PSL age 0, and PSL age 1+).

## 2.2 | Environmental data

We used two environmental data sets to examine the relationship between ocean climate and forage fish length and condition index: the basin-scale Pacific Decadal Oscillation (PDO) index and the local sea surface temperature (SST) over a ~4° latitude × 4° longitude area around each island site, encompassing most of the foraging range of the birds. The PDO is defined as the first empirical orthogonal function of detrended SST poleward of 20°N and represents an index of ocean temperature–climate variability in the North Pacific basin (Mantua & Hare, 2002). Monthly time series were averaged to produce the seasonal values winter: January–March (JFM); spring: April–June (AMJ); and summer: July–September (JAS). Each SST and PDO season was compared to annually averaged length and condition for each size/age class of fish separately to avoid model overfitting due to SST autocorrelation and because lagged relationships with the fish sampled in June–August may indicate different mechanisms of environmental influence. SST data were obtained from the Hadley Center HadISST1 reconstructed data set (<https://www.metoffice.gov.uk/hadobs/hadisst/>), which provides monthly 1-degree data for the study period (1993–2016). We selected a "coastal" box (52–56°N, 136–132°W) for St. Lazaria and a "Gulf of Alaska" box (54–58°N, 144–138°W) for Middleton (Figure 1), over which we averaged SST values to obtain one local SST index for each site. Seasonal SST time series did not have significant linear trends

at any site, while quadratic regressions showed significant ( $p < 0.05$ ) fits for spring (April–June) in both regions. This was largely due to the SST anomalies in 2014–2016 associated with the marine heat wave (Figure S5; Di Lorenzo & Mantua, 2016). When the 2015–2016 values were removed, these quadratic regressions were not significant ( $p > 0.05$ ), and therefore, we did not detrend the SST or PDO data.

Modeling the relationships between fish indices (length, *K* condition index) and ocean climate time series (SST, PDO) was done using weighted multiple regressions. Each regression included one fish index as a dependent variable and only one seasonal ocean climate index at a time as the independent variable. We tested linear and quadratic (2<sup>nd</sup>-degree polynomial) values of the ocean climate indices. The models were first evaluated by discarding any non-significant models using a *p*-value ( $p < 0.05$ ) criterion. In no case were the quadratic models significant (Tables S5 and S6). From the resulting significant models, we selected the best model for each fish index and each climate index, but across all seasons, based on the lowest Akaike information criterion (AIC) statistic. Secondly, we verified the selection of the best model by examination of the  $R^2$  value; a higher  $R^2$  value indicated greater explanatory power.


## 3 | RESULTS

### 3.1 | Length–frequency distributions

Due to the timing of sampling corresponding to the May–July spawning period for capelin, most capelin sampled by rhinoceros auklets were age 1 and older fish (hereafter age 1+), though a few small, probably age-0, fish were also sampled (Figure 2a,b; also see Naumenko, 1966; Pahlke, 1985). Owing to the few age-0 fish in our samples, we did not stratify capelin by size/age class for analyses. In contrast, a distinct bimodal structure for length was evident for PSL (Figure 2c,d). Typically, PSL < 100 mm in length during summer are assumed to be age-0 fish, whereas fish ≥100 mm are assumed to be primarily age 1+ (Arimitsu, Piatt, Heflin, Biela, & Schoen, 2018; Cunningham et al., 2018; Robards, Piatt, & Rose, 1999). It is possible, however, that in years of anomalous environmental conditions, PSL growth diminishes sufficiently such that some age-1 fish are misclassified as age 0 (von Biela et al., 2019). Since our data set included large samples of both smaller and larger PSL, we stratified analyses by these size/assumed age classifications to examine differences between these groups. There were significant differences in average length of fish sampled between regions (capelin:  $t = -12.26$ ,  $p < 0.0001$ ; PSL:  $t = 4.60$ ,  $p < 0.0001$ ), with larger fish for both species (PSL age 1+) found at Middleton in the northern GoA (Figure S1).

### 3.2 | Fulton's *K* index of body condition

Condition indices of capelin and age-0 PSL were significantly higher in the eastern GoA (St. Lazaria) in comparison with the northern region (Middleton); however, there was no significant difference in *K* between regions for age-1+ PSL (Table 2). There was significant inter-annual variability in fish condition for capelin and age-1+ PSL at



**FIGURE 2** Length–frequency histograms for capelin at (a) Middleton and (b) St. Lazaria islands, and Pacific sand lance at (c) Middleton and (d) St. Lazaria islands

Species	Site	N	Mean	SE	t	p ( T  >  t )
Capelin	Middleton	2,340	0.478	0.0036	-4.714	<0.0001
	St. Lazaria	1,139	0.503	0.0020		
PSL age 0	Middleton	1,055	0.267	0.0030	-7.132	<0.0001
	St. Lazaria	497	0.300	0.0022		
PSL age 1+	Middleton	876	0.329	0.0035	1.684	0.092
	St. Lazaria	451	0.320	0.0019		

Abbreviation: PSL, Pacific sand lance.

both sites and age-0 PSL at Middleton (Table 3, Figures 3 and S2). For capelin and PSL age 1+, we found that anomalies oscillated between positive and negative over the course of the time series in cycles of 4–5 years. Age-0 PSL showed less interannual variability in general (Table 3). There was a significant effect of sample date for capelin and both PSL age classes at Middleton, and PSL age 1+ at St. Lazaria, but not for capelin and PSL age 0 at St. Lazaria Island (Table 3). In all cases, the regression coefficient for sampling date was negative,

indicating a slight decline in condition over the course of the summertime sampling period (Figure S3).

### 3.3 | Relationships with environment

Capelin length was inversely related to spring (AMJ) SST and PDO at Middleton (Table 4, Figure 4). Spring SST was a better predictor of capelin length than the PDO, with 52% of the variation in length

**TABLE 2** Results of unpaired two-sided t test to test for differences in mean condition between each site by species and size/age class

**TABLE 3** Results of ANCOVA on body condition (Fulton's *K*) by year and sampling date for capelin and Pacific sand lance (PSL) at each site, 1993–2016

Species	Site	N fish	Variable	df	F	p > F	Coefficient
Capelin	Middleton	2,351	Model	12	9.81	<0.001	-0.0013
			Year	11	9.98	<0.001	
			Sample date	1	13.83	<0.001	
Capelin	St. Lazaria	1,688	Model	21	7.25	<0.001	-0.0001
			Year	20	7.55	<0.001	
			Sample date	1	0.55	0.457	
PSL age 0	Middleton	2,649	Model	12	7.64	<0.001	-0.0003
			Year	11	7.82	<0.001	
			Sample date	1	4.80	0.029	
PSL age 0	St. Lazaria	924	Model	16	1.16	0.294	-0.0003
			Year	15	1.22	0.248	
			Sample date	1	0.14	0.706	
PSL age 1+	Middleton	1,551	Model	13	3.93	<0.001	-0.0010
			Year	12	3.51	<0.001	
			Sample date	1	8.64	0.003	
PSL age 1+	St. Lazaria	1,315	Model	18	9.50	<0.001	-0.0006
			Year	17	9.84	<0.001	
			Sample date	1	18.68	<0.001	

explained by SST (Table 4). No relationship between capelin length and SST or PDO was established at St. Lazaria. At both Middleton and St. Lazaria islands, capelin body condition was inversely related to SST and the PDO, explaining between 42% and 68% of the variation in condition (Table 5, Figure 5). The summer (JAS) SST effect was more pronounced at Middleton, and spring (AMJ) temperature (SST and PDO) was the most important at St. Lazaria.

Environmental relationships for PSL varied by size/age class. Winter (JFM) SST and PDO explained 41%–46% of the variance in the length of age-0 PSL at Middleton, and the relationships were positive, in accordance with predictions (Table 4, Figure 4). However, no relationships were found for length of age-0 PSL at St. Lazaria. Spring (AMJ) SST and PDO explained 23%–26% of the variance in the length of age-1+ PSL at Middleton, but these relationships were negative, in contrast to our predictions (Table 4, Figure 4). Summer (JAS) PDO explained 25% of the variance in the body condition of age-0 PSL at St. Lazaria, but no relationships were found with Middleton data (Table 5, Figure 5). For age-1+ PSL, winter (JFM) SST accounted for 36% of the variance at Middleton, but no relationships were found for St. Lazaria (Table 5).


## 4 | DISCUSSION

Based on previous work that showed apparent habitat affinities with opposing responses by capelin and PSL to cool versus warm ocean temperatures (Sydeman, Piatt, et al., 2017), we predicted that capelin length and body condition would show negative relationships with both SST and the PDO, whereas the relationships for PSL length and body condition would be positive. We found support for our

hypothesis for capelin; however, temperature effects on the length and condition of PSL depended on their size/assumed age classes. For age-0 PSL, our findings suggest a positive relationship between length and winter SST, supporting our hypothesis, which may reflect earlier hatch dates and/or higher metabolic and growth rates (using length as a proxy) under warmer winter ocean conditions.

Assuming growth is slower for age-1+ PSL than age-0 PSL, the decrease in PSL age-1+ body condition with increasing temperatures could be related to poorer feeding conditions under warmer ocean conditions (Brodeur, Hunsicker, Hann, & Miller, 2019; Daly & Brodeur, 2015). This mechanism could potentially affect both age-1+ PSL and capelin. In support of this idea, von Biela et al. (2019) found that age-1 PSL experienced unusually low growth in 2015–2016 during the marine heat wave in the region (see Figure S1c; Di Lorenzo & Mantua, 2016). Slow growth and low abundance in sandeel populations in the North Sea were related to poor feeding conditions (MacDonald, Speirs, Greenstreet, Boulcott, & Heath, 2019), and it is reasonable to assume similar effects during the Northeast Pacific marine heat wave. Moreover, our selected models almost always included a lag of at least one season to the summer season of fish sampling. As noted in other studies (Byrd, Schmutz, & Renner, 2008; Shultz, Piatt, Harding, Kettle, & Pelt, 2009; Sydeman, Thompson, et al., 2017; Zador, Hunt, TenBrink, & Aydin, 2013), these relationships indicate that ocean climate in seasons prior to the summer growing period has notable effects, presumably operating through changes in production at lower trophic levels.

A key assumption of this study is that the samples provided by rhinoceros auklets are representative of the size and age classes of these forage fish in the environment. Differences in the lengths of capelin and PSL by region and year may be related to the



**FIGURE 3** Average annual body condition (Fulton's  $K$ ) anomalies over time at Middleton (black bars) and St. Lazaria (gray bars) for (a) capelin, (b) age-0 Pacific sand lance (PSL), and (c) age-1+ PSL. Dashed lines indicate  $\pm 1$  standard deviation

availability of different size fish in the foraging habitat surrounding the islands where sampling took place. Recent work has shown that seabirds and other predators sample species in proportion to their availability (Bertram & Kaiser, 1993; Piatt et al., 2018), and tagging studies indicate that seabirds sample PSL where net sampling methods underperform (Cunningham et al., 2018). Less is known, however, about whether seabirds sample size of prey in proportion to size availability. This same assumption would presumably also apply to any sampling device used, including net sampling techniques.

#### 4.1 | Length–frequency distributions and life history characteristics

In the GoA, capelin spawn inshore during May–July, and coastal areas may provide nursery grounds for larval fish (Arimitsu et al., 2008; Brown, 2002; Brown et al., 2002). In comparison, PSL spawn in shallow water during an extended period in fall and winter and peak hatching occurs in February–March (Doyle, Picquelle, Mier,

Spillane, & Bond, 2009; Robards, Piatt, et al., 1999). Sand lance begin spawning at roughly 20 months (age 1), and most spawn at age 2 or 3, while smaller percentages survive to age 4–6 (Robards, Piatt, et al., 1999).

Since spawning occurred during the previous fall/winter, age-0 PSL are of suitable size for seabird predation in the June–August sampling period. In contrast, due to the timing of the capelin spawning season, few age-0 capelin would be of the size suitable for sampling by the birds during summer, and correspondingly, we see few age-0 capelin in our samples. Indeed, the timing of capelin spawning may have a stronger effect on sampling at Middleton, which is roughly 100 km offshore of the nearest known spawning location (Arimitsu et al., 2018). On the other hand, St. Lazaria is coastal, so larval/juvenile capelin may be more abundant in this region. While we did not specifically analyze age (e.g., using otoliths) at either site, the bimodal distribution corresponding with small (<100 mm) and larger ( $\geq 100$  mm) classes of PSL generally indicates that both age-0 and age-1+ fish were taken, and in similar amounts. In Prince William Sound in 2016, however, von Biela et al. (2019) found that age-0 and

**TABLE 4** Best models based on AIC (lowest) and  $R^2$  (highest) from regressions of the PDO and regional SST by season on average annual body size (length, mm)

Species	Site	Model				Term	$\beta$	SE of $\beta$	t-stat	p-value	Figure
		AIC	$R^2$	df	p-value						
Capelin	Middleton	y = 1 + SST_AMJ				Intercept	109.360	1.123	97.38	<0.0001	4a
		242.0	0.52	20	<0.0005	SST_AMJ	-8.090	1.743	-4.64	<0.0005	
		y = 1 + PDO_AMJ				Intercept	107.440	1.380	77.71	<0.0001	4b
		250.0	0.29	20	<0.01	PDO_AMJ	-4.770	1.680	-2.85	<0.01	
	St. Lazaria	None									
	PSL age 0	Middleton	y = 1 + SST_JFM				Intercept	79.660	1.151	69.23	<0.0001
269.0			0.41	22	<0.001	SST_JFM	4.960	1.276	3.89	<0.001	
y = 1 + PDO_JFM				Intercept	79.980	1.032	77.53	<0.0001	4d		
267.0			0.46	22	<0.0005	PDO_JFM	3.550	0.825		4.30	<0.0005
St. Lazaria		None									
PSL age 1+		Middleton	y = 1 + SST_AMJ				Intercept	123.180	1.911	69.16	<0.0001
	297.0		0.26	22	0.011	SST_AMJ	-6.150	2.225	-2.76	0.011	
	y = 1 + PDO_AMJ				Intercept	131.400	1.832	71.72	<0.0001	4f	
	292		0.23	22	0.018	PDO_AMJ	-4.640	1.820	-2.55		0.018
	St. Lazaria	None									

Abbreviations: AMJ, spring, April–June; JAS, summer, July–September; JFM, winter, January–March; PSL, Pacific sand lance.

age-1 PSL lengths overlapped, but this was during the intense marine heat wave in the region. Potential bias in our ability to correctly classify age class from length is a possible issue in our study, which included these years. However, the 2014–2016 marine heat wave in the North Pacific has been described as unprecedented (Di Lorenzo & Mantua, 2016), meaning that the data from Prince William Sound during these years are probably anomalous. Otherwise, such extreme length-at-age anomalies are rare in *Ammodytes* (Macer, 1966; Wanless et al., 2018).


## 4.2 | Seasonal variability in condition indices

The body condition of capelin and PSL varied seasonally and depending on the date of sampling. Two main factors influence seasonal variability in condition. First, the condition of females of both capelin and PSL varies with the stage of reproductive maturity because gravid females have higher mass/length ratio and energy density than males (Carscadden & Frank, 2002; Montevecchi & Piatt, 1984; Robards, Anthony, et al., 1999). This can create a substantial difference because egg masses can increase the weight of gravid female capelin by up to 30% (Montevecchi & Piatt, 1984). Thus, a sample containing both males and females could produce a slightly biased (higher) condition index if a large proportion of the sample

was gravid females. This could have affected our findings for capelin because sampling of fish (June–August) partially overlaps with the spawning period of capelin (May–July) in Alaska (Arimitsu et al., 2008). It should not have affected our PSL samples, however, since they typically spawn later in the year (Robards, Anthony, et al., 1999). Second, seasonal variability in body mass may be related to life history strategies for lipid storage. Capelin store lipid in the fall and use it slowly over winter (Montevecchi & Piatt, 1984), and then more rapidly as they migrate from offshore to nearshore for spawning in the spring. Lipid content is lowest prior to and during spawning (Buren et al., 2014; Montevecchi & Piatt, 1984). By August, juveniles and surviving adults begin to quickly regain lipid reserves. PSL exhibit a similar seasonal cycle, except that peak lipid storage occurs in June–August and stored fat declines steadily until they spawn in the fall (Robards, Piatt, & Rose, 2002). Thus, for both capelin and PSL, the seasonal variability in timing of spawning and lipid storage probably contributes to the annual variability in body condition described here.

## 4.3 | Regional and interannual variation in condition

Condition values for capelin and age-0 PSL were generally higher in the eastern GoA (St. Lazaria Island) than in the northern GoA




**FIGURE 4** Predicted (line) and observed (asterisks) values of average annual length (mm) based on best models (see Table 4 for model results) for environmental parameters. Dashed lines reflect 95% confidence intervals for predictions. (a) Capelin at Middleton by spring SST, (b) capelin at Middleton by spring PDO, (c) Pacific sand lance (PSL) age 0 at Middleton by winter SST, (d) PSL age 0 at Middleton by winter PDO, (e) PSL age 1+ at Middleton by spring SST, and (f) PSL age 1+ at Middleton by spring PDO. No model was selected for any of the fish at St. Lazaria [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

**TABLE 5** Best models based on AIC (lowest) and  $R^2$  (highest) from regressions of the PDO and regional SST by season on average annual body condition (Fulton's  $K$ )

Species	Site	Model					Term	$\beta$	SE of $\beta$	t-stat	p-value	Figure
		AIC	$R^2$	df	p-value							
Capelin	Middleton	y = 1 + SST_JAS					Intercept	0.48	0.009	54.27	<0.0001	5a
		14.7	0.42	10	0.023	SST_JAS						
	St. Lazaria	y = 1 + SST_AMJ					Intercept	0.513	0.004	131.24	<0.0001	5b
		-17.3	0.61	19	<0.0001	SST_AMJ						
		y = 1 + PDO_AMJ					Intercept	0.503	0.003	148.9	<0.0001	5c
		-21.1	0.68	19	<0.0001	PDO_AMJ						
PSL age 0	Middleton	None					Intercept	0.325	0.011	28.91	<0.0001	5d
	St. Lazaria	y = 1 + PDO_JAS										
PSL age 1+	Middleton	y = 1 + SST_JFM					Intercept	0.330	0.006	51.72	<0.0001	5e
		-1.26	0.36	11	0.029	SST_JFM						
	St. Lazaria	None										

Abbreviations: AMJ, spring, April–June; JAS, summer, July–September; JFM, winter, January–March; PSL, Pacific sand lance.

**FIGURE 5** Predicted (line) and observed (asterisks) values of average annual body condition based on best models (see Table 5 for model results) for environmental parameters. Dashed lines reflect 95% confidence intervals for predictions. (a) Capelin at Middleton by summer SST, (b) capelin at St. Lazaria by spring SST, (c) capelin at St. Lazaria by spring PDO, (d) Pacific sand lance (PSL) age 0 at St. Lazaria by summer PDO, and (e) PSL age 1+ at Middleton by winter SST. No model was selected for PSL age 0 at Middleton or PSL age 1+ at St. Lazaria [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



(Middleton Island); no differences between the sites were found for age-1+ PSL (Table 2). Though the differences between these sites were small, there are a number of factors that may explain them, including environmental factors and/or variation in sample size. St. Lazaria Island is a coastal site located <20 km offshore (56°N), whereas Middleton is situated at the edge of the continental shelf, 75 km offshore in the northern GoA (59°N). Both islands and their biota may be influenced by the Alaska Current, formed as the northern bifurcation of the North Pacific Current (Hollowed & Wooster, 1992). Local physical oceanographic processes, however, differ substantially. St. Lazaria is located closer to the bifurcation streamline of the North Pacific Current (Cummins & Freeland, 2007), and in a coastal domain where freshwater runoff and downwelling are important factors to ecosystem productivity. In contrast, Middleton is located in more oceanic waters and is less influenced by coastal enrichments. Conditions at both locations are influenced by the PDO, although larger influence has been hypothesized for the northern GoA than on the eastern side (Lluch-Cota, Wooster, & Hare, 2001). Due to this oceanographic variation, it is possible that the coastal waters near St. Lazaria are, in general, more productive than the offshore waters near Middleton. Indeed, Waite and Mueter (2013) showed that satellite-derived chlorophyll-*a* indices in May–July, prior to and overlapping the beginning of our sampling period, were generally higher in southeast GoA

than in the northern-central GoA. Whether this variation in phytoplankton standing stocks links to variation among higher trophic levels is currently unknown.

Interannual variability in body condition for capelin and PSL was significant in almost all cases, with an exception of age-0 PSL at St. Lazaria. Young-of-the-year fish typically distribute energy toward growth rather than lipid storage (Burril, Biela, Hillgruber, & Zimmerman, 2018; Robards, Anthony, et al., 1999), so this finding is not surprising (Brosset et al., 2015). In another study overlapping with ours in study period, the energy density of age-0 PSL in Prince William Sound did not vary in 2012–2016, despite large variation in physical conditions between years (von Biela et al., 2019). Similarly, there was relatively low interannual variability in condition of age-0 PSL at Middleton, though condition for these fish was much lower in 1 year, 2008, following a very cold and late spring (Figure 3; Hatch, 2013).

#### 4.4 | Temperature and variation in condition

Models including SST or the PDO explained a relatively large percentage (25%–68%) of the variance in capelin and PSL body condition in the GoA. SST and PDO had negative effects on capelin and age-1+ PSL body condition, indicating lower body condition with warming for these larger and older fish. The effects of SST and the

PDO, however, were positive for age-0 PSL, suggesting that warmer ocean temperatures are important for younger, developing fish, provided that food for them is not limited. It was summertime SST or PDO that related positively to body condition of age-0 PSL, whereas wintertime ocean climate was related negatively to body condition of older fish.

In contrast, Carscadden and Frank (2002) found no correlation between capelin body condition and temperature in Newfoundland; however, they did find high capelin condition during the 1980s followed by a decade of low condition in the 1990s. Later studies revealed that this decadal-scale change in condition coincided with a regime shift in which capelin and cod (*Gadus morhua*) populations both crashed after 1990 and remained extremely low until after the mid-2000s (Buren et al., 2014, 2019; Rose & Rowe, 2015). In this region, zooplankton stocks also crashed during the 1990s, so the food supply for capelin may have been limited (Carscadden & Frank, 2002). Buren et al. (2014) found evidence that capelin population biomass was indirectly affected by temperature via effects on timing of spring sea ice retreat and a corresponding change in phenology of primary and secondary productivity. Elsewhere, evidence suggests that changes in zooplankton prey communities from rich- to poor-quality species, coupled with an increase in fish metabolic rate under warming, were responsible for a reduction in body condition of juvenile salmonids off Oregon (Brodeur et al., 2019; Daly & Brodeur, 2015). Similarly, long-term studies of anchovy and sardine in the Mediterranean Sea suggest that while temperature changes may be associated with changes in body condition (especially in sardine), the most consistent and significant predictor of body condition was meso-zooplankton concentration (Brosset et al., 2015). In summary, evidence suggests that marked, persistent changes in body condition indicate changes in bottom-up processes that affect food webs and these are often coincident with or driven by temperature changes that may or may not be associated with regime shifts.

That said, there were no obvious signs in our study of effects from the 1997–1998 El Niño event, or the unusual climate conditions that took place in 2005 that affected marine food webs around the northeast Pacific (Brodeur et al., 2006). On the other hand, the persistence of the 2014–2016 marine heat wave (Bond, Cronin, Freeland, & Mantua, 2015; Di Lorenzo & Mantua, 2016) was evident in the length–frequency distribution of PSL at Middleton, which was largely unimodal in 2015 and 2016 and consisted primarily of the small/age-0 class (Figure S4). Additionally, body condition of PSL age 1+ at Middleton in 2016 was the lowest value of the time series, and nearly the same as the average condition for the small PSL (~0.29). von Biela et al. (2019) reported that in Prince William Sound, the energy density of age-1 PSL began to decline in 2015, and the 2015 PSL year class experienced little growth over the winter of 2015–2016. This anomalous growth was an 89% decline in whole-body energy for age-1 PSL compared to cooler years, which may have occurred because of poor feeding conditions, particularly during the 2015 summer/fall PSL growth season.

## 5 | CONCLUSION

We found species-specific, regional, and interannual variation in forage fish length and body condition, suggesting that climate variability in the North Pacific can have substantial impacts on prey species morphometrics, and hence energy density, in the GoA. Changes in condition of key prey species may have cascading effects in marine food webs, and this will be a topic of future analyses. For example, condition of capelin and PSL was among the lowest of the time series in 2015–2016 and may have contributed to the large-scale die-off of seabirds and unusual mortality of marine mammals in the region during the fall and winter of 2015–2016 (Walsh et al., 2018). This study thus provides another step toward a greater understanding of variation in forage fish length and condition due to climate variability in the GoA.

## ACKNOWLEDGEMENTS

We thank the North Pacific Research Board (Project 1706), National Fish and Wildlife Foundation, and U.S. Geological Survey Alaska Science Center for supporting this analysis. Heather Renner, Brie Drummond, Sarah Laske, and two anonymous reviewers provided thorough and very thoughtful feedback on the manuscript that further improved it, and we express our gratitude to them as well. We also sincerely appreciate the efforts of dozens of field biologists who collected data on Middleton and St. Lazaria islands over the past several decades. Any use of trade, firm, or product names is for descriptive purposes only and does not constitute endorsement by the U.S. government.

## CONFLICT OF INTEREST

The authors of this paper have no conflict of interest to declare.

## AUTHOR CONTRIBUTIONS

JFP, SAH, and MGR contributed data to the study. SAT, WJS, and MGR were responsible for data analysis. SAT, WJS, MLA, MGR, and JFP wrote, edited, and revised the manuscript.

## DATA ACCESSIBILITY

The Pacific Decadal Oscillation index is available at [https://oceanview.pfeg.noaa.gov/erddap/tabledap/cciea\\_OC\\_PDO.html](https://oceanview.pfeg.noaa.gov/erddap/tabledap/cciea_OC_PDO.html). Sea surface temperature data were obtained from the Hadley Center HadISST1 reconstructed data set (<https://www.metoffice.gov.uk/hadobs/hadisst/>). The forage fish data that support the findings of this study are available on request from the corresponding author. Those data are not publicly available due to privacy or ethical restrictions.

## ORCID

Sarah Ann Thompson  <https://orcid.org/0000-0003-1667-0853>

William J. Sydeman  <https://orcid.org/0000-0003-1902-4654>

## REFERENCES

- Abookire, A. A., & Piatt, J. F. (2005). Oceanographic conditions structure forage fishes into lipid-rich and lipid-poor communities in lower Cook Inlet, Alaska, USA. *Marine Ecology Progress Series*, 287, 229–240. <https://doi.org/10.3354/meps287229>
- Alaska Maritime National Wildlife Refuge (AMNWR). (2018). *Standardized protocols for annual seabird monitoring camps at Aiktak, Buldir, Chowiet, St. George, St. Lazaria and St. Paul islands and Cape Lisburne in the Alaska Maritime National Wildlife Refuge in 2018*. U.S. Fish and Wildlife Service Report, AMNWR 2018/04. Homer, AK.
- Alverson, D. L. (1992). A review of commercial fisheries and the Steller sea lion *Eumetopias jubatus*: The conflict arena. *Reviews in Aquatic Sciences*, 6, 203–256.
- Anthony, J. A., Roby, D. D., & Turco, K. R. (2000). Lipid content and energy density of forage fishes from the northern Gulf of Alaska. *Journal of Experimental Marine Biology & Ecology*, 248, 53–78. [https://doi.org/10.1016/S0022-0981\(00\)00159-3](https://doi.org/10.1016/S0022-0981(00)00159-3)
- Arimitsu, M. L., Piatt, J. F., Heflin, B., von Biela, V. R., & Schoen, S. K. (2018). *Monitoring long-term changes in forage fish distribution, abundance, and body condition*. Anchorage, AK: U.S. Geological Survey, Alaska Science Center.
- Arimitsu, M. L., Piatt, J. F., Litzow, M. A., Abookire, A. A., Romano, M. D., & Robards, M. D. (2008). Distribution and spawning dynamics of capelin (*Mallotus villosus*) in Glacier Bay, Alaska: A cold water refugium. *Fisheries Oceanography*, 17, 137–146. <https://doi.org/10.1111/j.1365-2419.2008.00470.x>
- Baker, M. R., Matta, M. E., Beaulieu, M., Paris, N., Huber, S., Graham, O. J., ... O'Neill, M. R. (2019). Intra-seasonal and inter-annual patterns in the demographics of sand lance and response to environmental drivers in the North Pacific. *Marine Ecology Progress Series*, 617–618, 221–244. <https://doi.org/10.3354/meps12897>
- Bertram, D. F., & Kaiser, G. W. (1993). Rhinoceros auklets (*Cerorhinca monocerata*) nestling diet may gauge Pacific sand lance (*Ammodytes hexapterus*) recruitment. *Canadian Journal of Fisheries and Aquatic Sciences*, 50, 1908–1915.
- Bond, N. A., Cronin, M. F., Freeland, H., & Mantua, N. (2015). Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophysical Research Letters*, 42, 3414–3420. <https://doi.org/10.1002/2015gl063306>
- Brodeur, R. D., Hunsicker, M. E., Hann, A., & Miller, T. W. (2019). Effects of warming ocean conditions on feeding ecology of small pelagic fishes in a coastal upwelling ecosystem: A shift to gelatinous food sources. *Marine Ecology Progress Series*, 617–618, 149–163. <https://doi.org/10.3354/meps12497>
- Brodeur, R. D., Ralston, S., Emmett, R. L., Trudel, M., Auth, T. D., & Phillips, A. J. (2006). Anomalous pelagic nekton abundance, distribution, and apparent recruitment in the northern California Current in 2004 and 2005. *Geophysical Research Letters*, 33, L22S08. <https://doi.org/10.1029/2006GL026614>
- Brodeur, R. D., & Ware, D. M. (1992). Long-term variability in zooplankton biomass in the subarctic Pacific Ocean. *Fisheries Oceanography*, 1, 32–38. <https://doi.org/10.1111/j.1365-2419.1992.tb00023.x>
- Brosset, P., Ménard, F., Fromentin, J.-M., Bonhommeau, S., Ulses, C., Bourdeix, J.-H., ... Sarau, C. (2015). Influence of environmental variability and age on the body condition of small pelagic fish in the Gulf of Lions. *Marine Ecology Progress Series*, 529, 219–231. <https://doi.org/10.3354/meps11275>
- Brown, E. D. (2002). Life history, distribution, and size structure of Pacific capelin in Prince William Sound and the northern Gulf of Alaska. *ICES Journal of Marine Science*, 59, 983–996. <https://doi.org/10.1006/jmsc.2002.1281>
- Brown, E. D., Churnside, J. H., Collins, R. L., Veenstra, T., Wilson, J. J., & Abnett, K. (2002). Remote sensing of capelin and other biological features in the North Pacific using LIDAR and video technology. *ICES Journal of Marine Science*, 59, 1120–1130. <https://doi.org/10.1006/jmsc.2002.1282>
- Buren, A. D., Koen-Alonso, M., Pepin, P., Mowbray, F. K., Nakashima, B., Stenson, G. B., ... Montevecchi, W. A. (2014). Bottom-up regulation of capelin, a keystone forage species. *PLoS ONE*, 9, e87589. <https://doi.org/10.1371/journal.pone.0087589>
- Buren, A. D., Murphy, H. M., Adamack, A. T., Davoren, G. K., Koen-Alonso, M., Montevecchi, W. A., ... Varkey, D. (2019). The collapse and continued low productivity of a keystone forage fish species. *Marine Ecology Progress Series*, 616, 155–170. <https://doi.org/10.3354/meps12924>
- Burger, A. E., Wilson, R. P., Garnier, D., & Wilson, M.-P.-T. (1993). Diving depths, diet, and underwater foraging of rhinoceros auklets in British Columbia. *Canadian Journal of Zoology*, 71, 2528–2540. <https://doi.org/10.1139/z93-346>
- Burril, S. E., von Biela, V. R., Hillgruber, N., & Zimmerman, C. E. (2018). Energy allocation and feeding ecology of juvenile chum salmon (*Oncorhynchus keta*) during transition from freshwater to salt-water. *Polar Biology*, 41, 1447–1461. <https://doi.org/10.1007/s00300-018-2297-2>
- Byrd, G. V. (2007). Seabird monitoring on Alaska Maritime National Wildlife Refuge. In R. Brewer (Ed.), *Community-based coastal observing in Alaska: Aleutian Life Forum 2006* (pp. 39–45). Fairbanks, AK: Alaska Sea Grant.
- Byrd, G. V., Schmutz, J. A., & Renner, H. M. (2008). Contrasting population trends of piscivorous seabirds in the Pribilof Islands: A 30-year perspective. *Deep Sea Research Part II: Topical Studies in Oceanography*, 55, 1846–1855. <https://doi.org/10.1016/j.dsr2.2008.04.004>
- Carscadden, J. E., & Frank, K. T. (2002). Temporal variability in the condition factors of Newfoundland capelin (*Mallotus villosus*) during the past two decades. *ICES Journal of Marine Science*, 59, 950–958. <https://doi.org/10.1006/jmsc.2002.1234>
- Carscadden, J. E., Montevecchi, W. A., Davoren, G. K., & Nakashima, B. S. (2002). Trophic relationships among capelin (*Mallotus villosus*) and seabirds in a changing ecosystem. *ICES Journal of Marine Science*, 59, 1027–1033. <https://doi.org/10.1006/jmsc.2002.1235>
- Crawford, R. J. M., Sydesman, W. J., Thompson, S. A., Sherley, R. B., & Makhado, A. B. (2019). Food habits of an endangered seabird indicate recent poor availability of abundant forage resources. *ICES Journal of Marine Science*, in press. <https://doi.org/10.1093/icesjms/fsz081>
- Cummins, P. F., & Freeland, H. J. (2007). Variability of the North Pacific current and its bifurcation. *Progress in Oceanography*, 75, 253–265. <https://doi.org/10.1016/j.pocean.2007.08.006>
- Cunningham, J. T., Elliott, K. H., Cottenie, K., Hatch, S. A., & Jacobs, S. R. (2018). Individual foraging location, but not dietary, specialization: Implications for rhinoceros auklets as samplers of forage fish. *Marine Ecology Progress Series*, 605, 225–240. <https://doi.org/10.3354/meps12761>
- Daly, E. A., & Brodeur, R. D. (2015). Warming ocean conditions relate to increased trophic requirements of threatened and endangered salmon. *PLoS ONE*, 10, 1–23. <https://doi.org/10.1371/journal.pone.0144066>
- Davoren, G. K., & Montevecchi, W. A. (2003). Signals from seabirds indicate changing biology of capelin stocks. *Marine Ecology Progress Series*, 258, 253–261. <https://doi.org/10.3354/meps258253>
- Di Lorenzo, E., & Mantua, N. (2016). Multi-year persistence of the 2014/15 North Pacific marine heatwave. *Nature Climate Change*, 6, 1042–1048. <https://doi.org/10.1038/nclimate3082>
- Doyle, M. J., Picquelle, S. J., Mier, K. L., Spillane, M. C., & Bond, N. A. (2009). Larval fish abundance and physical forcing in the Gulf of Alaska, 1981–2003. *Progress in Oceanography*, 80, 163–189. <https://doi.org/10.1016/j.pocean.2009.03.002>
- Friesen, V. L., Baker, A. J., & Piatt, J. F. (1996). Phylogenetic relationships within the Alcidae (Charadriiformes: Aves) inferred from total

- molecular evidence. *Molecular Biology and Evolution*, 13, 359–367. <https://doi.org/10.1093/oxfordjournals.molbev.a025595>
- Hatch, S. A. (2013). Kittiwake diets and chick production signal a 2008 regime shift in the Northeast Pacific. *Marine Ecology Progress Series*, 477, 271–284. <https://doi.org/10.3354/meps10161>
- Hatch, S. A., & Sanger, G. A. (1992). Puffins as samplers of juvenile pollock and other forage fish in the Gulf of Alaska. *Marine Ecology Progress Series*, 80, 1–14. <https://doi.org/10.3354/meps080001>
- Hollowed, A. B., & Wooster, W. S. (1992). Variability of winter ocean conditions and strong year classes of Northeast Pacific groundfish. *ICES Marine Science Symposium*, 195, 433–444.
- IPCC (2013). *Climate Change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, NY: Cambridge University Press.
- Kuroki, M., Kato, A., Watanuki, Y., Niizuma, Y., Takahashi, A., & Naito, Y. (2003). Diving behavior of an epipelagically feeding alcid, the rhinoceros auklet (*Cerorhinca monocerata*). *Canadian Journal of Zoology*, 81, 1249–1256. <https://doi.org/10.1139/z03-112>
- Litz, M. N. C., Brodeur, R. D., Emmett, R. L., Heppell, S. S., Rasmussen, R. S., O'Higgins, L., & Morris, M. S. (2010). Effects of variable oceanographic conditions on forage fish lipid content and fatty acid composition in the northern California Current. *Marine Ecology Progress Series*, 405, 71–85. <https://doi.org/10.3354/meps08479>
- Lluch-Cota, D. B., Wooster, W. S., & Hare, S. R. (2001). Sea surface temperature variability in coastal areas of the Northeastern Pacific related to the El Niño-Southern Oscillation and the Pacific Decadal Oscillation. *Geophysical Research Letters*, 28, 2029–2032. <https://doi.org/10.1029/2000GL012429>
- Logerwell, E. A., & Schaufler, L. E. (2005). New data on proximate composition and energy density of Steller sea lion (*Eumetopias jubatus*) prey fills seasonal and geographic gaps in existing information. *Aquatic Mammals*, 31, 62–82. <https://doi.org/10.1578/AM.31.1.2005.62>
- Logerwell, E. A., Stabeno, P. J., Wilson, C. D., & Hollowed, A. B. (2007). The effect of oceanographic variability and interspecific competition on juvenile pollock (*Theragra chalcogramma*) and capelin (*Mallotus villosus*) distributions on the Gulf of Alaska shelf. *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 2849–2868. <https://doi.org/10.1016/j.dsr2.2007.08.008>
- MacDonald, A., Speirs, D. C., Greenstreet, S. P. R., Boulcott, P., & Heath, M. R. (2019). Trends in sandeel growth and abundance off the East Coast of Scotland. *Frontiers in Marine Science*, 6, 201. <https://doi.org/10.3389/fmars.2019.00201>
- Macer, C. (1966). Sand eels (Ammodytidae) in the south-western North Sea; their biology and fishery. *Fishery Investigations Series II*, 24, 1–55.
- Mantua, N. J., & Hare, S. R. (2002). The Pacific Decadal Oscillation. *Journal of Oceanography*, 58, 35–44.
- Mills, K. L., Laidig, T., Ralston, S., & Sydeman, W. J. (2007). Diets of top predators indicate pelagic juvenile rockfish (*Sebastes* spp.) abundance in the California Current System. *Fisheries Oceanography*, 16, 273–283. <https://doi.org/10.1111/j.1365-2419.2006.00429.x>
- Montevecchi, W. A., & Piatt, J. (1984). Composition and energy contents of mature inshore spawning capelin *Mallotus villosus* implications for sea bird predators. *Comparative Biochemistry & Physiology A*, 78, 15–20. [https://doi.org/10.1016/0300-9629\(84\)90084-7](https://doi.org/10.1016/0300-9629(84)90084-7)
- Nash, R. D. M., Valencia, A. H., & Geffen, A. J. (2006). The origin of Fulton's condition factor – setting the record straight. *Fisheries*, 31, 236–238.
- Naumenko, E. (1966). Distribution, biological condition, and abundance of capelin (*Mallotus villosus socialis*) in the Bering Sea. In O. A. Mathisen, & K. O. Coyle (Eds.), *Ecology of the Bering Sea: A review of Russian literature* (pp. 237–256). Alaska Sea Grant Report No. 96–01. Fairbanks, AK: Alaska Sea Grant College Program, University of Alaska Fairbanks.
- Osterblom, H., Olsson, O., Blenckner, T., & Furness, R. W. (2008). Junk-food in marine ecosystems. *Oikos*, 117, 967–977. <https://doi.org/10.1111/j.2008.0030-1299.1650.x>
- Pahlke, K. A. (1985). *Preliminary studies of capelin (Mallotus villosus) in Alaskan waters. Informational Leaflet No. 250*. Juneau, AK: Alaska Department of Fish and Game.
- Pangle, K. L., & Sutton, T. M. (2005). Temporal changes in the relationship between condition indices and proximate composition of juvenile *Coregonus artedii*. *Journal of Fish Biology*, 66, 1060–1072. <https://doi.org/10.1111/j.1095-8649.2005.00660.x>
- Payne, S. A., Johnson, B. A., & Otto, R. S. (1999). Proximate composition of somenorth-easternPacific forage fish species. *Fisheries Oceanography*, 8, 159–177. <https://doi.org/10.1046/j.1365-2419.1999.00097.x>
- Piatt, J. F., Arimitsu, M. L., Sydeman, W. J., Thompson, S. A., Renner, H., Zador, S., ... Williams, J. (2018). Biogeography of pelagic food webs in the North Pacific. *Fisheries Oceanography*, 27, 366–380. <https://doi.org/10.1111/fog.12258>
- Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of California*, 191, 1–382.
- Robards, M. D., Anthony, J. A., Rose, G. A., & Piatt, J. F. (1999). Changes in proximate composition and somatic energy content for Pacific sand lance (*Ammodytes hexapterus*) from Kachemak Bay, Alaska relative to maturity and season. *Journal of Experimental Marine Biology and Ecology*, 242, 245–258. [https://doi.org/10.1016/s0022-0981\(99\)00102-1](https://doi.org/10.1016/s0022-0981(99)00102-1)
- Robards, M. D., Piatt, J. F., & Rose, G. A. (1999). Maturation, fecundity, and intertidal spawning of Pacific sand lance in the northern Gulf of Alaska. *Journal of Fish Biology*, 54, 1050–1068. <https://doi.org/10.1006/jfbi.1999.0941>
- Robards, M. D., Piatt, J. F., & Rose, G. A. (2002). Growth and abundance of Pacific sand lance, *Ammodytes hexapterus*, under differing oceanographic regimes. *Environmental Biology of Fishes*, 64, 429–441. <https://doi.org/10.1023/A:1016151224357>
- Rose, G. A., & O'Driscoll, R. L. (2002). Capelin are good for cod: Can the northern stock rebuild without them? *ICES Journal of Marine Science*, 59, 1018–1026. <https://doi.org/10.1006/jmsc.2002.1252>
- Rose, G. A., & Rowe, S. (2015). Northern cod comeback. *Canadian Journal of Fisheries and Aquatic Sciences*, 72, 1789–1798. <https://doi.org/10.1139/cjfas-2015-0346>
- Rosen, D. A. S., & Trites, A. W. (2000). Pollock and the decline of Steller sea lions: Testing the junk-food hypothesis. *Canadian Journal of Zoology*, 78, 1243–1250. <https://doi.org/10.1139/z00-060>
- Schoen, S. K., Piatt, J. F., Arimitsu, M. L., Heflin, B. M., Madison, E. N., Drew, G. S., ... DeGange, A. R. (2018). Avian predator buffers against variability in marine habitats with flexible foraging behavior. *Marine Biology*, 165, 47. <https://doi.org/10.1007/s00227-018-3304-4>
- Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, 1, 401–406. <https://doi.org/10.1038/nclimate1259>
- Shultz, M. T., Piatt, J. F., Harding, A. M. A., Kettle, A. B., & Van Pelt, T. I. (2009). Timing of breeding and reproductive performance in murrelets and kittiwakes reflect mismatched seasonal prey dynamics. *Marine Ecology Progress Series*, 393, 247–258. <https://doi.org/10.3354/meps08136>
- Stabeno, P. J., Bell, S., Cheng, W., Danielson, S., Kachel, N. B., & Mordy, C. W. (2016). Long-term observations of Alaska Coastal Current in the northern Gulf of Alaska. *Deep Sea Research Part II: Topical Studies in Oceanography*, 132, 24–40. <https://doi.org/10.1016/j.dsr2.2015.12.016>
- Stabeno, P. J., Bond, N. A., Hermann, A. J., Kachel, N. B., Mordy, C. W., & Overland, J. E. (2004). Meteorology and oceanography of the Northern Gulf of Alaska. *Continental Shelf Research*, 24, 859–897. <https://doi.org/10.1016/j.csr.2004.02.007>

- Suryan, R. M., Irons, D. B., Kaufman, M., Benson, J., Jodice, P. G. R., Roby, D. D., & Brown, E. D. (2002). Short-term fluctuations in forage fish availability and the effect on prey selection and brood-rearing in the black-legged kittiwake *Rissa tridactyla*. *Marine Ecology Progress Series*, 236, 273–287. <https://doi.org/10.3354/meps236273>
- Sydeman, W. J., Piatt, J. F., Thompson, S. A., Garcia-Reyes, M., Hatch, S. A., Arimitsu, M. L., ... Renner, H. M. (2017). Puffins reveal contrasting relationships between forage fish and ocean climate in the North Pacific. *Fisheries Oceanography*, 26, 379–395. <https://doi.org/10.1111/fog.12204>
- Sydeman, W. J., Thompson, S. A., Piatt, J. F., Garcia-Reyes, M., Zador, S. G., Williams, J. C., ... Renner, H. M. (2017). Regionalizing indicators for marine ecosystems: Bering Sea-Aleutian Island seabirds, climate, and competitors. *Ecological Indicators*, 78, 458–469. <https://doi.org/10.1016/j.ecolind.2017.03.013>
- Thayer, J. A., Bertram, D. F., Hatch, S. A., Hipfner, M. J., Slater, L., Sydeman, W. J., & Watanuki, Y. (2008). Forage fish of the Pacific Rim as revealed by diet of a piscivorous seabird: Synchrony and relationships with sea surface temperature. *Canadian Journal of Fisheries and Aquatic Sciences*, 65, 1610–1622. <https://doi.org/10.1139/F08-076>
- Trites, A. W., & Donnelly, C. P. (2003). The decline of Steller sea lions *Eumetopias jubatus* in Alaska: A review of the nutritional stress hypothesis. *Mammal Review*, 33, 3–28. <https://doi.org/10.1046/j.1365-2907.2003.00009.x>
- Van Pelt, T. I., Piatt, J. F., Lance, B. K., & Roby, D. D. (1997). Proximate composition and energy density of some North Pacific forage fishes. *Comparative Biochemistry and Physiology, Part A*, 118, 1393–1398. [https://doi.org/10.1016/S0300-9629\(97\)00240-5](https://doi.org/10.1016/S0300-9629(97)00240-5)
- Vollenweider, J. J., Heintz, R. A., Schaufler, L., & Bradshaw, R. (2011). Seasonal cycles in whole-body proximate composition and energy content of forage fish vary with water depth. *Marine Biology*, 158, 413–427. <https://doi.org/10.1007/s00227-010-1569-3>
- von Biela, V. R., Arimitsu, M. L., Piatt, J. F., Heflin, B., Schoen, S., Trowbridge, J., & Clawson, C. (2019). Extreme reduction in condition of a key forage fish during the Pacific marine heatwave of 2014–2016. *Marine Ecology Progress Series*, 613, 171–182. <https://doi.org/10.3354/meps12891>
- Waite, J. N., & Mueter, F. J. (2013). Spatial and temporal variability of chlorophyll-a concentrations in the coastal Gulf of Alaska, 1998–2011, using cloud-free reconstructions of SeaWiFS and MODIS-Aqua data. *Progress in Oceanography*, 116, 179–192. <https://doi.org/10.1016/j.pocean.2013.07.006>
- Walsh, J. E., Thoman, R. L., Bhatt, U. S., Bieniek, P. A., Brettschneider, B., Brubaker, M., ... Partain, J. (2018). The high latitude marine heat wave of 2016 and its impacts on Alaska. In: Explaining extreme events of 2016 from a climate perspective *Bulletin of the American Meteorological Society*, 99(1), S39–S43. <https://doi.org/10.1175/BAMS-D-17-0105.1>
- Wanless, S., Harris, M. P., Newell, M. A., Speakman, J. R., & Daunt, F. (2018). Community-wide decline in the occurrence of lesser sand-eels *Ammodytes marinus* in seabird chick diets at a North Sea colony. *Marine Ecology Progress Series*, 600, 193–206. <https://doi.org/10.3354/meps12679>
- Wanless, S., Harris, M. P., Redman, P., & Speakman, J. R. (2005). Low energy values of fish as a probable cause of a major seabird breeding failure in the North Sea. *Marine Ecology Progress Series*, 294, 1–8. <https://doi.org/10.3354/meps294001>
- Zador, S., Hunt, G. L. Jr, TenBrink, T., & Aydin, K. (2013). Combined seabird indices show lagged relationships between environmental conditions and breeding activity. *Marine Ecology Progress Series*, 485, 245–258. <https://doi.org/10.3354/meps10336>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Thompson SA, García-Reyes M, Sydeman WJ, Arimitsu ML, Hatch SA, Piatt JF. Effects of ocean climate on the length and condition of forage fish in the Gulf of Alaska. *Fish Oceanogr.* 2019;28:658–671. <https://doi.org/10.1111/fog.12443>